





CQIP

Cancer Quality Improvement Program Newton-Wellesley Hospital


6141530

Newton, MA



Annual Report 2014

Colon Cancer: Adjuvant chemotherapy for lymph node positive colon cancer



	My Program	My State (MA)	Region (New			All CoC Programs
Performance Rate	100 %	90.1 %	88.8 %	88.8 %	87.7 %	86.4 %


Adjuvant chemotherapy is considered or administered within 4 months (120 days) of diagnosis for patients under the age of 80 with AJCC Stage III (lymph node positive) colon cancer. (CP3R data as of 12/12/2014)







Colon Cancer: At least 12 regional lymph nodes removed and pathologically examined for resected colon cancer



	My Program	My State (MA)	My Census Region (New England)		My CoC Program Type (COMP)	All CoC Programs
Performance Rate	96.9 %	90.1 %	89.4 %	89.4 %	87.8 %	88.1 %


At least 12 regional lymph nodes are removed and pathologically examined for resected colon cancer. (CP3R data as of 12/12/2014)







Stage Distribution - Colon Cancer Diagnosed in 2012, NWH vs. All CoC



	0		II	III	IV	NA	UNK
My Facility	0 % (n=0)	18.64 % (n=11)	25.42 % (n=15)	30.51 % (n=18)	22.03 % (n=13)	0 % (n=0)	3.39 % (n=2)
All CoC	5.77 % (n=4272)	19.39 % (n=14350)	24.26 % (n=17954)	24.72 % (n=18293)	19.84 % (n=14682)	0.12 % (n=87)	5.89 % (n=4357)





